

Significant contributions of Albrecht's A term to non-resonant Raman scattering processes

Zu-Yong Gong,^{1,2} Guangjun Tian,² Sai Duan,^{2, a)} and Yi Luo^{1,2}

¹⁾Hefei National Laboratory for Physical Science at the Microscale, Department of Chemical Physics, School of Chemistry and Materials Science, University of Science and Technology of China, Hefei, 230026 Anhui, P. R. China.

²⁾Department of Theoretical Chemistry and Biology, School of Biotechnology, Royal Institute of Technology, S-106 91 Stockholm, Sweden.

(Dated: 10 August 2015)

The Raman intensity can be well described by the famous Albrecht equation that consists of A and B terms. It is well known that the contribution from Albrecht's A term can be neglected without loss of accuracy for far off-resonant Raman scattering processes. However, as demonstrated in this study, we have found that this widely accepted long-standing assumption fails drastically for totally symmetric vibration modes of molecules in general off-resonant Raman scattering. Perturbed first principles calculations for water molecule show that strong constructive interference between the A and B terms occurs for the Raman intensity of the symmetric O-H stretching mode, which can account for about 40% of the total intensity. Meanwhile, a minor destructive interference is found for the angle bending mode. The state to state mapping between the Albrecht's theory and the perturbation theory allows us to verify the accuracy of the widely employed perturbation method for the dynamic/resonant Raman intensities. The model calculations rationalized from water molecule with the bending mode show that the perturbation method is a good approximation only when the absolute energy difference between the first excited state and the incident light is more than five times of the vibrational energy in ground state.

I. INTRODUCTION

Raman spectroscopy¹ is one of the standard vibrational spectroscopic tools that has been extensively applied in different fields. The intensity of the Raman scattering can be generally treated by Albrecht's Raman theory², in which the polarizability is expanded into two terms, *i.e.* the famous Albrecht's A and B terms^{2,3}

$$\alpha_{pq} = A_{pq} + B_{pq}, \quad (1)$$

where p and q represent Cartesian coordinates. If both initial and final electronic states belong to the ground state, they can be written as^{2,3}

$$\begin{aligned} A_{pq} &= \frac{1}{\hbar} \sum_{r \neq g} \sum_{v^r=0}^{\infty} \frac{p_{e^g e^r}^0 q_{e^r e^g}^0}{\omega_{e^r v^r: e^g v^i} - \omega} \langle v^f | v^r \rangle \langle v^r | v^i \rangle \\ &+ \frac{1}{\hbar} \sum_{r \neq g} \sum_{v^r=0}^{\infty} \frac{p_{e^g e^r}^0 q_{e^r e^g}^0}{\omega_{e^r v^r: e^g v^f} + \omega} \langle v^f | v^r \rangle \langle v^r | v^i \rangle \\ B_{pq} &= \frac{1}{\hbar} \sum_{r \neq g} \sum_{v^r=0}^{\infty} \frac{p_{e^g e^r}^{k,0} q_{e^r e^g}^0}{\omega_{e^r v^r: e^g v^i} - \omega} \langle v^f | Q_k | v^r \rangle \langle v^r | v^i \rangle \\ &+ \frac{1}{\hbar} \sum_{r \neq g} \sum_{v^r=0}^{\infty} \frac{p_{e^g e^r}^0 q_{e^r e^g}^{k,0}}{\omega_{e^r v^r: e^g v^f} + \omega} \langle v^f | Q_k | v^r \rangle \langle v^r | v^i \rangle \\ &+ \frac{1}{\hbar} \sum_{r \neq g} \sum_{v^r=0}^{\infty} \frac{p_{e^g e^r}^0 q_{e^r e^g}^{k,0}}{\omega_{e^r v^r: e^g v^i} - \omega} \langle v^f | v^r \rangle \langle v^r | Q_k | v^i \rangle \\ &+ \frac{1}{\hbar} \sum_{r \neq g} \sum_{v^r=0}^{\infty} \frac{p_{e^g e^r}^{k,0} q_{e^r e^g}^0}{\omega_{e^r v^r: e^g v^f} + \omega} \langle v^f | v^r \rangle \langle v^r | Q_k | v^i \rangle, \end{aligned} \quad (2)$$

where $\hbar$ is the reduced Planck's constant, $|e^g\rangle$ and $|e^r\rangle$ are electronic ground and excited states, $|v^i\rangle$ and $|v^f\rangle$ are the initial and final vibrational states of $|e^g\rangle$ associated with frequency ω^g , $|v^r\rangle$ is the vibrational state of $|e^r\rangle$ associated with frequency ω^r , ω is the frequency of the incident light, $\omega_{e^r v^r: e^g v^i}$ is the frequency difference between $|e^r\rangle|v^r\rangle$ and $|e^g\rangle|v^i\rangle$, $p_{e^g e^r}^0$ is the transition dipole moment between $|e^g\rangle$ and $|e^r\rangle$ at equilibrium geometry (Q_0), $p_{e^g e^r}^{k,0}$ is the derivative of $p_{e^g e^r}^0$ with respect to specific normal mode Q_k . Here we should emphasize that *all* modes could contribute as intermediate states in the Raman processes. In other words, $|v^r\rangle$ represents all possible combination of all modes, *i.e.*, $|v^r\rangle = |v_1^r v_2^r \cdots v_N^r\rangle$, where N is the number of vibrational modes. In some cases, the denominator in Eq. 2 would be written in terms of vertical excitation energy ΔE_{rg} , which obeys the relationship $\Delta E_{rg} = \hbar\omega_{e^r: e^g} + \lambda$. Here λ is the reorganization energy.⁴

The Albrecht's theory is commonly used to study the resonant Raman scattering,⁵ although it is a general theory for all Raman processes³. It is well known that, for strongly dipole-allowed transitions, A term is dominant in the resonant Raman spectra.^{5,6} It is noted that A term does not contribute to non totally symmetric modes.⁷ On the other hand, for weakly dipole-allowed transitions, both the A and B terms could significantly contribute to the Raman scattering of totally symmetric modes. In these cases, the relative magnitude of them could only be determined by quantum chemical calculations.⁸⁻¹¹ It is also noted that the effects of higher-order terms were also discussed in recent studies.¹¹

For *non*-resonant Raman scattering processes, we focus on the zero-frequency limit, *i.e.*, the ω is neglected.

^{a)}Electronic mail: said@theochem.kth.se

In the traditional treatment of the Albrecht's theory for *non-resonant* situations, it has been often assumed that $\omega_{e^r v^r; e^g v^i} \approx \omega_{e^r; e^g}$.² As a result, the denominator in Eq. 2 is independent of the vibrational state $|v^r\rangle$. Therefore, the summation of v^r could be calculated prior. Because of the completeness of $|v^r\rangle$ and orthonormality between $|v^i\rangle$ and $|v^f\rangle$, the prior summation will result in the A term only responsible for the Rayleigh scattering and vanishing for *non-resonant* Raman scattering.² Nowadays, this argument has been widely accepted in the theory of *non-resonant* Raman scattering.^{3,7,12,13} However, the assumption is apparently too strong. If we only consider the fact that $\Delta E_{rg} \gg \hbar\omega_{v^r} - \hbar\omega_{v^i} - \lambda$, in the framework of harmonic approximation and expanding the denominator by Taylor series, the component related to an *arbitrary* excited state $|e^r\rangle$ in Eq. 2 can be rewritten as

$$\begin{aligned}
 A_{pq}^r &= \frac{2p_{e^g e^r}^0 q_{e^r e^g}^0}{\Delta E_{rg}} \langle v^f | v^i \rangle \\
 &+ \frac{p_{e^g e^r}^0 q_{e^r e^g}^0}{\Delta E_{rg}^2} [(1 + v^i + v^f) \hbar\omega^g + 2\lambda - \hbar\omega^r] \langle v^f | v^i \rangle \\
 &- \frac{2p_{e^g e^r}^0 q_{e^r e^g}^0}{\Delta E_{rg}^2} \sum_{v^r=0}^{\infty} \hbar v^r \omega^r \langle v^f | v^r \rangle \langle v^r | v^i \rangle \\
 B_{pq}^r &= \frac{2(p_{e^g e^r}^{k,0} q_{e^r e^g}^0 + p_{e^g e^r}^0 q_{e^r e^g}^{k,0})}{\Delta E_{rg}} \langle v^f | Q_k | v^i \rangle \\
 &+ \mathcal{O}\left(\frac{\hbar\omega_{v^r} - \hbar\omega_{v^i} - \lambda}{\Delta E_{rg}}\right).
 \end{aligned} \tag{3}$$

Here we should emphasize the multimode nature of ω^g and ω^r . For instance, we have the expression of $\hbar v^r \omega^r = \sum_{k=1}^N \hbar v_k^r \omega_k^r$. The schematic drawings for a single vibrational mode of all definitions could be found in Fig. 1, where ω_k and ω'_k are the vibrational frequencies related to $|e^g\rangle$ and $|e^r\rangle$, respectively. The summation A_{pq}^r over all $|e^r\rangle$ would return to A_{pq} . If we consider the Stokes shift for the fundamental frequency, *i.e.* $v^i = 0$ and $v^f = 1$, the first two terms in A_{pq}^r vanish due to the orthogonality of the vibrational wavefunctions in the ground state. However, the last term survives and should contribute to the *non-resonant* Raman intensity.

The finite difference method¹⁴ and coupled-perturbed method¹⁵ have also been used to compute the intensity of *non-resonant* Raman spectra by directly differentiating the electronic polarizabilities, *i.e.*^{3,16}

$$\alpha_{pq} = \langle v^f | \alpha_{e,pq} | v^i \rangle, \tag{4}$$

where¹⁷

$$\alpha_{e,pq} = \sum_{r \neq g} \frac{p_{e^g e^r} q_{e^r e^g}}{\Delta E_{rg} \pm \hbar\omega}. \tag{5}$$

As a result, the component related to state $|e^r\rangle$ in Eq. 4




FIG. 1. Schematic drawing of the single displaced harmonic model used to study the contribution of the A term to the Raman intensity. Q_0 (ω_k) and Q_r (ω'_k) are the equilibrium geometries (vibrational frequencies) of ground state $|e^g\rangle$ and excited state $|e^r\rangle$, respectively. All other symbols can be found in Eq. 3.

can be rewritten as

$$\begin{aligned}
 \alpha_{pq}^r &= \frac{2p_{e^g e^r}^0 q_{e^r e^g}^0}{\Delta E_{rg}} \langle v^f | v^i \rangle \\
 &- \frac{2p_{e^g e^r}^0 q_{e^r e^g}^0}{\Delta E_{rg}^2} \frac{\partial \Delta E_{rg}}{\partial Q_k} \langle v^f | Q_k | v^i \rangle \\
 &+ \frac{2(p_{e^g e^r}^{k,0} q_{e^r e^g}^0 + p_{e^g e^r}^0 q_{e^r e^g}^{k,0})}{\Delta E_{rg}} \langle v^f | Q_k | v^i \rangle.
 \end{aligned} \tag{6}$$

Again, the zero-frequency limit is also applied in Eq. 6 and the summation α_{pq}^r would become α_{pq} . In both cases, the Raman cross section can be calculated from the polarizabilities in Eqs. 1 and 4.¹⁶

In this work, we comprehensively investigate the relationship between Eqs. 6 and 3. Comparing Eqs. 6 and 3, we can immediately find that the B_{pq}^r term corresponds to the last term in α_{pq}^r . Thus, the focus of current study is the relationship for A term. We first address this issue in the model system and show that the A term indeed contributes to final results for *non-resonant* conditions. Then, the A term in time frame is briefly discussed. Finally, we take water monomer as a realistic example to show the importance of including the A term for final results.

II. MODEL SYSTEM

The first hint for the corresponding relationship for A term between Eqs. 6 and 3 can be found in the case of $\partial\Delta E_{rg}/\partial Q_k = 0$. If we consider that both potential energy surfaces (PESs) of $|e^g\rangle$ and $|e^r\rangle$ are harmonic, this case is equivalent to the situation that the displacement between the two PESs is 0 ($\Delta Q = 0$). We can immediately obtain that the second term in α_{pq}^r is 0. On the other hand, because of the no shift in PESs, either $\langle 1|v^r\rangle$ or $\langle v^r|0\rangle$ will be 0 due to the parity symmetry.⁷ As a result, the last A_{pq}^r term is also 0 for Raman scattering. Hence, one could notice that the last term in A_{pq}^r may correspond to the second term in α_{pq}^r . It is well known that the case can be realized in the non-totally symmetric vibrations.⁷ For such kind of vibrations, there always exists at least one symmetric operator that makes the geometries of $Q_0 \pm \delta Q_k$ to be identity and then $\partial\Delta E_{rg}/\partial Q_k = 0$. Thus, the reason of A term has no contribution to the *non*-resonant Raman intensity of non-totally symmetric vibrations is the orthogonality of vibrational wavefunctions between ground and excited states rather than the orthogonality of two vibrational wavefunctions in the ground state. The former reason is exactly the same as that for the selection rule in resonant Raman processes.^{5,7,12} It is worth to note that this conclusion can also be generalized to resonant Raman intensities as well as general PESs (such as double well¹⁸) of excited states.

To further confirm the corresponding relationship between the last term in A_{pq}^r and the second term in α_{pq}^r , the condition of $\partial\Delta E_{rg}/\partial Q_k \neq 0$ should be enforced. In the framework of linear coupling model (LCM)¹⁹ where it assumes $\omega_k = \omega'_k$, both terms can be calculated analytically and give the same result, *i.e.*,

$$\frac{\sqrt{2\hbar\omega_k^3}p_{eg}^0q_{er}^0\Delta Q}{\Delta E_{rg}^2}. \quad (7)$$

When $\Delta Q \geq 0$, this result is equal to $2p_{eg}^0q_{er}^0\hbar\omega_k\sqrt{S}/\Delta E_{rg}^2$, where S is the Huang-Rhys factor²⁰. We should emphasize that the current treatment is consistent with Ting's work²¹ which could be traced back to Shorygin's treatment²² in 1947. However, Ting's work was restricted to the LCM scheme and the vibronic coupling (the B term) has not been included.²³ Moreover, Ting's algorithm was only widely applied in the two-state model²⁴⁻²⁶ and intrinsically pre-resonant conditions^{24,25,27}. In off-resonance conditions, Ting's A term was usually considered to be very small and negligible.^{23,28}

When going beyond the LCM, with the help of the general sum rules for Franck-Condon integrals,²⁹ we could obtain that the last term in A_{pq}^r and the second term in α_{pq}^r both equal to

$$\frac{\sqrt{2\hbar\omega_k^2}p_{eg}^0q_{er}^0\Delta Q}{\sqrt{\omega_k}\Delta E_{rg}^2}. \quad (8)$$

We have further performed numerical calculations for more general condition of $\partial\Delta E_{rg}/\partial Q_k \neq 0$. The considered displaced harmonic model is shown in Fig. 1. For example, in the case of $\omega_k = 1600 \text{ cm}^{-1}$, $\omega'_k = 1800 \text{ cm}^{-1}$, and $\Delta Q_k = Q_r - Q_0 = 10 \text{ a.u.}$, both terms give the same value of 0.0055704918 in the unit of $2p_{eg}^0q_{er}^0\hbar/\Delta E_{rg}^2$. The analytical and perfect numerical agreement not only confirm the equivalence of the two terms but also show that, for modes with $\partial\Delta E_{rg}/\partial Q_k \neq 0$, the A term can have non-zero contribution to the *non*-resonant Raman intensity. Note that, here the corresponding S is around 0.4,³⁰ which is reasonable for realistic molecules. Because the $|e^r\rangle$ is arbitrary in above discussions, a state to state relationship between Albrecht's theory and perturbation method is thus established for *non*-resonant Raman spectrum. It is worth to mention that, in the original work of Albrecht, a similar relationship has also been discussed. However, it only focused on the case of near-resonance conditions (see Eq. 22 in Ref. 2).

III. TIME-DEPENDENT FRAME

Time-dependent frame of Raman theory has the advantage of avoiding the sum-over-states of $|v^r\rangle$ in Albrecht's theory.^{18,23,31-33} Instead, the dynamics of the wave packet given by the initial vibrational wave function times the corresponding electronic transition dipole on the excited states is involved.²³ Mathematically, the equivalence of these two representations could be readily proven by a simple half-Fourier transform.^{23,31} In time frame, the "dynamic" and "static" terms are equivalent to the Albrecht's A and B terms, respectively. Here, the "dynamic" term arises from wave packet propagation on the Born-Oppenheimer surfaces, while, the "static" term comes from the coordinate dependence of the electronic transition dipole.²³ In other words, the first and second terms in the Taylor expansion of transition dipole, *i.e.* $p_{er}^0 = p_{er}^0 + p_{er}^{k,0}Q_k$, associate with the "dynamic" and "static" terms, respectively.

In *non*-resonant conditions, the short time approximation holds. Thus, in the zero-frequency limit, the time average polarizability associated with $|e^r\rangle$ could be calculated as^{23,34}

$$\bar{\alpha}_{pq}^r = \frac{2\hbar}{T} \int_0^T \alpha_{pq}^r(s) ds, \quad (9)$$

where the per-factor 2 arises from the zero-frequency limit, $T = 2\pi\hbar/|E_{av} - E_{egv^i}|$, and

$$\alpha_{pq}^r(s) = \int_0^s dt \exp\left(\frac{-iE_{egv^i}t}{\hbar}\right) \langle \phi_f | \exp(i\mathcal{H}_r t/\hbar) | \phi_i \rangle. \quad (10)$$

Here $\mathcal{H}_r$ is the vibrational Hamiltonian for $|e^r\rangle$, the damping factor is omitted, $\langle \phi_f | = \langle v^f | p_{eg}^r$, $|\phi_i\rangle = q_{er}^0 |v^i\rangle$, and E_{av} is the average energy on $|e^r\rangle$. If the assumption^{23,35}

$$\langle \phi_f | \mathcal{H}_r - E_{av} | \phi_i \rangle = 0 \quad (11)$$




FIG. 2. Three vibrational modes of water molecule.

is adopted, the “dynamic” term in Eq. 10 would be

$$\alpha_{pq}^{r,D}(s) = \int_0^s dt \exp\left(\frac{i\Delta E t}{\hbar}\right) p_{eg}^0 q_{eg}^0 \langle v^f | v^i \rangle, \quad (12)$$

where $\Delta E = E_{av} - E_{eg} v^i$. As a result, the “dynamic” term (the A term) of the Stokes peaks at fundamental frequencies also vanishes in the time frame due to the orthogonality of the vibrational wavefunctions in the ground state. On the other hand, the “static” terms survives and its value in the first order of Q_k is exactly equal to B term in Eq. 3 if we considered $\Delta E = \Delta E_{rg}$ (see Appendix B in Ref. 23 for details). This conclusion is consistent with that in Ref. 23, where the emphasis for far from resonance is solely on the “static” term.

The reason of the vanishing of “dynamic” term in *non*-resonance in the time frame would be easily located as the assumption of Eq. 11, which is equivalent to the assumption of $\omega_{ev^r v^r; eg v^i} \approx \omega_{ev^r; eg}$ in the energy frame. Both assumptions neglect the vibrational contributions in excited states. To obtain the correct result, in the time frame, Eq. 9 should be calculated with the inclusion of vibrational contributions in $\mathcal{H}_r$, so that the contribution of the “dynamic” term (Albrecht’s A term) will be included in the total intensity.

IV. FIRST PRINCIPLES EXAMPLE: WATER MONOMER

Although the contribution of A term for *non*-resonant Raman intensities has been briefly touched,^{21–23} its magnitude for realistic molecules was still controversial^{25,28}. Therefore, a first principles example is highly desirable to address this issue. To the best of our knowledge, there were no previous examples that have investigated the magnitudes of both A and B terms at first principles level for *non*-resonant Raman intensities. In the following, we take water molecule as a realistic example to fully examine the contribution of the A term. Water molecule has three vibrational modes,³⁶ which are $\angle$ H-O-H bending (v_2), symmetric O-H stretching (v_1), and asymmetric O-H stretching modes (v_3), respectively. The vibrational modes of water are depicted in Fig. 2. All required excited states except core-hole excitations were calculated at time-dependent density functional theory level.³⁷ The finite difference method³⁸ was used to calculate the derivatives in α_{pq}^r . All calculated α_{pq}^r were summed up

TABLE I. Calculated Raman scattering factors for water (in the unit of $\text{\AA}^4/\text{amu}$) contain only the A term (A), the B term (B), or all terms (Tot.), as well as analytical results (Anal.). The experimental measurements (Expt.) extracted from Refs. 15, 40, and 41 also included for comparison.

Mode	Symmetry	A	B	Tot.	Anal.	Expt.
v_2	A_1	0.14	1.54	1.10	1.10	0.9 ± 0.2
v_1	A_1	11.66	48.32	104.94	104.54	108 ± 14
v_3	B_2	0.00	25.68	25.68	25.67	19.2 ± 2.1

over all singly excited states to obtain α_{pq} and finally Raman scattering factors were calculated by^{16,39}

$$S_k = 45a_k^2 + 7\gamma_k^2, \quad (13)$$

where

$$\begin{aligned} a_k &= \frac{1}{3} \{ (\alpha_{xx})_k + (\alpha_{yy})_k + (\alpha_{zz})_k \} \\ \gamma_k^2 &= \frac{1}{2} \left\{ [(\alpha_{xx})_k - (\alpha_{yy})_k]^2 \right. \\ &\quad + [(\alpha_{yy})_k - (\alpha_{zz})_k]^2 \\ &\quad + [(\alpha_{zz})_k - (\alpha_{xx})_k]^2 \\ &\quad \left. + 6 [(\alpha_{xy})_k^2 + (\alpha_{xz})_k^2 + (\alpha_{yz})_k^2] \right\}. \end{aligned} \quad (14)$$

Here the subscript “ k ” represents the individual vibrational mode. By using Eq. 6, we have calculated the scattering factors from three cases: only the A term (S_A), only the B term (S_B), and both A and B terms (S_{Tot}). With these scattering factors, the corresponding Raman intensities, *i.e.*, I_A , I_B and I_{Tot} , are calculated as^{3,16}

$$\begin{aligned} I &= \sum_k \frac{\pi^2}{\epsilon_0^2} (\tilde{\nu}_{\text{in}} - \tilde{\nu}_k)^4 \\ &\quad \times \frac{h}{8\pi^2 c \tilde{\nu}_k} \frac{S_k}{45} \frac{1}{1 - \exp(-hc\tilde{\nu}_k/k_B T)}, \end{aligned} \quad (15)$$

where ϵ_0 is the vacuum permittivity, $\tilde{\nu}_{\text{in}}$ and $\tilde{\nu}_k$ are the wave number of incident light and vibrational mode, respectively, h is the Planck’s constant, c is the speed of light, k_B is the Boltzmann constant, and T is the temperature. By definition, the interference contribution is calculated by $I_{\text{Tot}} - I_A - I_B$. For comparison, the analytical Raman scattering factors at the same density functional theory level were also computed.

All calculated scattering factors as well as experimental values^{40,41} are listed in Table I. As expected, the calculated S_{Tot} are identical to the analytical results and in good agreement with the experimental values^{15,40,41}. Here a slight overestimation for v_3 should be attributed to the approximate exchange-correlation functionals, since, at the CASSCF(10,10) level with the same basis set used here, the calculated value ($20.4 \text{ \AA}^4/\text{amu}$) is in the range of experimental observation.⁴² For the v_3 mode




FIG. 3. Calculated Raman intensities for water (black solid line). The contributions from the A term (red dotted), the B term (blue dotted), and the interference (black dotted) were also shown. The incident light and temperature were set to be 514.5 nm and 400 K followed by the experimental conditions,^{40,41} respectively. All spectra have been broadened by Lorentzian function with a full width at the half-maximum of 10 cm^{-1} .

that belongs to B_2 symmetry, the A term does not contribute to the total intensity, consistent with the discussion mentioned above. For the v_1 and v_2 modes, on the other hand, the contribution of the A term can no longer be neglected. For instance, the calculated S_B of v_1 is less than half of the analytical value. It is interesting to note that S_A can only account for about 10% of the analytical result and the summation of contributions from S_A and S_B terms can not provide the correct answer neither. On the other hand, a simple estimation shows that the value from $(\sqrt{S_A} + \sqrt{S_B})^2$ ($107 \text{ Å}^4/\text{amu}$) can reproduce the analytical result nicely. It indicates that a strong constructive interference between the A and B terms does exist for the v_1 mode and covers almost 40% of the total intensity. A minor destructive interference is identified for the v_2 mode. In this case, the simple summation of S_A and S_B gives a value larger than the final one. Contributions from different terms for the Raman intensities are also displayed in Fig. 3, which emphasizes the importance of the interference for the relative intensity. Notice that the first excitation energy of water is around 7 eV, which obviously obeys the condition of $\omega_{e^r v^r:eg} v^i \approx \omega_{e^r:eg}$. But even though, the contribution of the A term cannot be neglected. Overall, although the water monomer is a simple case, it is fully adequate to address our argument.

We notice that simple extension of Eq. 6 by replacing ΔE_{rg}^2 and ΔE_{rg} with $2[\Delta E_{rg} \pm \hbar(\omega + i\Gamma)]^2$ and $2[\Delta E_{rg} \pm \hbar(\omega + i\Gamma)]$, respectively,⁴³ was employed to calculate dynamic⁴⁴ and resonant Raman⁴⁵ intensities. Here the damping factor Γ represents the life time of the excited state. In addition, the same extension of the second term (the A term, labeled as $A_{pq}^{r,s}$ hereafter) in Eq. 6 has been used to explain the chemical enhancement of surface-enhanced Raman spectroscopy.^{46,47} However, the accuracy of such extension is questionable because,




FIG. 4. Relative error of simply extended A term in Eq. 6 respect to Eq. 2 at different energy of incident light for the model in Fig. 1. Here all parameters are rationalized from the first excited state of water molecule with the bending mode in LCM. Specifically, $\Delta Q = 1.35 \text{ a.u.}$, $\omega_k = \omega'_k = 1600 \text{ cm}^{-1}$, and $\omega_{e^r:eg} = 35 \omega_k$. Two Γ of 100 and 800 cm^{-1} were used. The insert figure shows the incident light around the resonant situation.

near the resonant condition, the Taylor expansion for denominator in Eq. 2 may fail and thus $|v^r\rangle\langle v^r|$ cannot be summed prior.⁴⁸

To verify the accuracy of this simplified extension of Eq. 6, we computed the relative error of $A_{pq}^{r,s}$ respect to Eq. 2 at different energy of incident light for the model system shown in Fig. 1. The result is depicted in Fig. 4. Here all parameters were rationalized from the first excited state of water molecule with the bending mode in LCM (see caption of Fig. 4 for details). Two typical Γ , *i.e.* small one of 100 cm^{-1} and large one of 800 cm^{-1} ,^{10,49} were used. Around the resonant situation, different Γ introduces different behaviour for the relative error. For instance, the maximum relative error for small Γ is much larger than that for large Γ . For the specific case, the maximum relative errors are even larger than 1400% and 130% for small and large Γ , respectively. In the region of away from resonant condition, the identical behaviour for relative errors with different Γ is observed in Fig. 4. Due to the uncertainty of experimental measurements as well as the limitation of the double harmonic approximation,⁵⁰ we used 15% as the threshold for the calculated $A_{pq}^{r,s}$. According to the model calculations, we have found that $A_{pq}^{r,s}$ is a good approximation when $|\omega_{e^r:eg} - \omega|$ is larger than $5 \omega_k$. We should emphasize that this value is mainly determined by ΔQ between two PESs. For example, when $\Delta Q = 10 \text{ a.u.}$, the range of accurate $A_{pq}^{r,s}$ is $|\omega_{e^r:eg} - \omega| > 10 \omega_k$. Apparently, when $\Delta Q = 0$, the parity symmetry will lead to zero A term both in Eq. 2 and Eq. 6.

V. CONCLUSION

We have clearly shown that the contributions from the Albrecht's A term are too large to be negligible for the *non*-resonant Raman intensities of totally symmetric fundamental vibrational modes, in contrast with the common wisdom of the field. The widely employed perturbation method is also found to be too crude for evaluating dynamic or resonant Raman spectra. Our findings are conceptually important for correctly understanding and modeling of Raman scattering processes under different conditions.

ACKNOWLEDGMENTS

This work was supported by the Ministry of Science and Technology of China (2010CB923300), the "Strategic Priority Research Program" of the Chinese Academy of Sciences (Grant XDB01020200), the Natural Science Foundation of China (21421063), and Swedish Research Council (VR). The Swedish National Infrastructure for Computing (SNIC) was acknowledged for computer time.

- ¹C. V. Raman and K. S. Krishnan, *Nature* **121**, 501 (1928).
- ²A. C. Albrecht, *J. Chem. Phys.* **34**, 1476 (1961).
- ³D. A. Long, *The Raman Effect: A Unified Treatment of the Theory of Raman Scattering by Molecules* (Wiley, Chichester New York, 2002).
- ⁴R. A. Marcus, *J. Chem. Phys.* **24**, 966 (1956).
- ⁵A. Myers Kelley, *J. Phys. Chem. A* **112**, 11975 (2008).
- ⁶D. W. Silverstein and L. Jensen, *J. Chem. Phys.* **136**, 064110 (2012).
- ⁷J. L. McHale, *Molecular Spectroscopy*, 1st ed. (Prentice Hall, Upper Saddle River, N.J., 1998).
- ⁸F. Santoro, C. Cappelli, and V. Barone, *J. Chem. Theory Comput.* **7**, 1824 (2011).
- ⁹D. W. Silverstein and L. Jensen, *J. Chem. Phys.* **136**, 064111 (2012).
- ¹⁰H. Ma, J. Liu, and W. Liang, *J. Chem. Theory Comput.* **8**, 4474 (2012).
- ¹¹F. Egidi, J. Bloino, C. Cappelli, and V. Barone, *J. Chem. Theory Comput.* **10**, 346 (2014).
- ¹²E. Smith and G. Dent, *Modern Raman Spectroscopy: A Practical Approach*, 1st ed. (John Wiley & Sons, Ltd, Hoboken, NJ, 2005).
- ¹³J. R. Lombardi and R. L. Birke, *J. Phys. Chem. C* **112**, 5605 (2008).
- ¹⁴I. G. John, C. B. Bacskey, and N. S. Hush, *Chem. Phys.* **38**, 319 (1979).
- ¹⁵R. D. Amos, *Chem. Phys. Lett.* **124**, 376 (1986).
- ¹⁶J. Neugebauer, M. Reiher, C. Kind, and B. A. Hess, *J. Comput. Chem.* **23**, 895 (2002).
- ¹⁷E. Merzbacher, *Quantum Mechanics*, 2nd ed. (Wiley, New York, 1970).
- ¹⁸S.-Y. Lee, *J. Chem. Educ.* **62**, 561 (1985).
- ¹⁹P. Macak, Y. Luo, and H. Ågren, *Chem. Phys. Lett.* **330**, 447 (2000).
- ²⁰K. Huang and A. Rhys, *Proc. R. Soc. Lond. A* **204**, 406 (1950).
- ²¹C.-H. Ting, *Spectrochim. Acta A* **24**, 1177 (1968).
- ²²P. P. Shorygin, *Zh. Fiz. Khim.* **12**, 1125 (1947).
- ²³S.-Y. Lee and E. J. Heller, *J. Chem. Phys.* **71**, 4777 (1979).
- ²⁴C. Castiglioni, M. Del Zoppo, and G. Zerbi, *Phys. Rev. B* **53**, 13319 (1996).
- ²⁵C. Castiglioni, M. Del Zoppo, and G. Zerbi, *Phys. Rev. B* **56**, 2275 (1997).
- ²⁶C.-C. Chou and B.-Y. Jin, *Theor. Chem. Acc.* **122**, 313 (2009).
- ²⁷E. S. Yeung, M. Heiling, and G. J. Small, *Spectrochim. Acta A* **31**, 1921 (1975).
- ²⁸D. M. Bishop and B. Kirtman, *Phys. Rev. B* **56**, 2273 (1997).
- ²⁹A. J. Sadlej, *Spectrochim. Acta A* **26**, 421 (1970), For facilitate obtaining current result, the vibrational Hamiltonian for $|e^r\rangle$ would be rewritten as $\mathcal{H}_r = \frac{1}{2}(p^2 + \omega_k'^2 Q'^2 - \hbar\omega_k')$, where, in the ground state coordinate, the mass-weighted coordinate on $|e^r\rangle$ could be expressed as $Q' = Q - \Delta Q$.
- ³⁰A. Alkauskas, J. L. Lyons, D. Steiauf, and C. G. Van de Walle, *Phys. Rev. Lett.* **109**, 267401 (2012).
- ³¹E. J. Heller, *Acc. Chem. Res.* **14**, 368 (1981).
- ³²E. J. Heller, R. Sundberg, and D. Tannor, *J. Phys. Chem.* **86**, 1822 (1982).
- ³³D. J. Tannor and E. J. Heller, *J. Chem. Phys.* **77**, 202 (1982).
- ³⁴Here the sign is consistent with Eq. 2 and the missed pre-factor $1/\hbar$ in Ref. 23 is added.
- ³⁵J. Tang and A. C. Albrecht, in *Raman Spectroscopy*, edited by H. A. Szymanski (Springer US, 1970) pp. 33–68.
- ³⁶F. Cotton, *Chemical Applications of Group Theory*, 3rd ed. (Wiley, New York, 1990).
- ³⁷The used functional and basis set were B3LYP implemented in Gaussian 09 and Sadlej pVTZ, respectively.
- ³⁸B. Fornberg, *Math. Comp.* **51**, 699 (1988).
- ³⁹E. B. Wilson Jr., J. C. Decius, and P. C. Cross, *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra* (Dover Publications, New York, 1980).
- ⁴⁰W. F. Murphy, *Mol. Phys.* **33**, 1701 (1977).
- ⁴¹W. F. Murphy, *Mol. Phys.* **36**, 727 (1978).
- ⁴²J. Neugebauer, M. Reiher, and B. A. Hess, *J. Chem. Phys.* **117**, 8623 (2002).
- ⁴³The opposite-sign convention for $1/\hbar$ is adopted, see Chapter A18 in Ref. 3.
- ⁴⁴C. Van Caillie and R. D. Amos, *Phys. Chem. Chem. Phys.* **2**, 2123 (2000).
- ⁴⁵D. Rappoport, S. Shim, and A. Aspuru-Guzik, *J. Phys. Chem. Lett.* **2**, 1254 (2011).
- ⁴⁶L. Jensen, C. M. Aikens, and G. C. Schatz, *Chem. Soc. Rev.* **37**, 1061 (2008).
- ⁴⁷A. T. Zayak, Y. S. Hu, H. Choo, J. Bokor, S. Cabrini, P. J. Schuck, and J. B. Neaton, *Phys. Rev. Lett.* **106**, 083003 (2011).
- ⁴⁸Although the sum-over-states in Eq. 5 is circumvented, the implicit prior summation for the $|v^r\rangle\langle v^r|$ is still included in the method in Ref. 44.
- ⁴⁹L. Zhao, L. Jensen, and G. C. Schatz, *J. Am. Chem. Soc.* **128**, 2911 (2006).
- ⁵⁰W. Koch and M. C. Holthausen, *A Chemist's Guide to Density Functional Theory*, 2nd ed. (Wiley-VCH Verlag GmbH, Weinheim New York, 2001).